

Bubbles and Quantum Tunnelling in Inflationary Cosmology

Stefano Ansoldi¹

*International Center for Relativistic Astrophysics (ICRA), Italy, and
Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Trieste, Italy, and
Dipartimento di Matematica e Informatica, Università degli Studi di Udine,
via delle Scienze 206, I-33100 Udine (UD), Italy [Mailing address]*

Abstract

We review a procedure to use semiclassical methods in the quantization of General Relativistic shells and apply these techniques in some simplified models of inflationary cosmology. Some interesting open issues are introduced and the relevance of their solution in the broader context of Quantum Gravity is discussed.

The interplay of the gravitational and quantum realms is a fundamental topic in the research landscape of the last decades and is still waiting for a consolidate answer. While waiting, it is sometimes also tempting to study simplified models, that are well known from the classical point of view and that can be turned into virtual laboratories to test our present level of understanding. Concerning quantum gravitational systems, it is safe to say that, if not most, certainly many of these models use *general relativistic shells*. To make a first, quick, contact with this interesting system, we will restrict to a highly symmetric case and consider two four-dimensional spherically symmetric spacetimes $\mathcal{S}_\pm$; let us also choose the coordinates $(t_\pm, r_\pm, \theta_\pm, \phi_\pm)$ which are static and adapted to the spherical symmetry, so that in both $\mathcal{S}_\pm$ the two metrics can be written as $g_{ab}^{(\pm)} = \text{diag}(g_{00}^{(\pm)}, g_{11}^{(\pm)}, g_{22}^{(\pm)}, g_{33}^{(\pm)}) = \text{diag}(-f_\pm(r_\pm), 1/f_\pm(r_\pm), r_\pm^2, r_\pm^2 \sin^2 \theta_\pm)$. Let us then consider the situation in which a part $\mathcal{M}_-$ of $\mathcal{S}_-$ and a part $\mathcal{M}_+$ of $\mathcal{S}_+$ are joined together across a timelike hypersurface Σ whose constant time slices are spherically symmetric, so that we obtain a new spherically symmetric spacetime $\mathcal{S} = \mathcal{M}_- \cup \Sigma \cup \mathcal{M}_+$. The dynamics of this system is described by Israel *junction conditions* [1], which, in the case we are considering, reduce to just one equation


$$R \left(\epsilon_+ \sqrt{\dot{R}^2 + f_+(R)} - \epsilon_- \sqrt{\dot{R}^2 - f_-(R)} \right) = M(R); \quad (1)$$

$M(R)$ is a function describing the matter content of Σ (i.e. it is related to the stress energy tensor of the infinitesimally thin matter-energy distribution which is *joining* $\mathcal{M}_-$ and $\mathcal{M}_+$); $\epsilon_\pm$ are the signs of the radicals which follow them: when $\epsilon_\pm$ are positive (resp. negative) it means that the normal to the shell (which by convention we choose directed from $\mathcal{M}_-$ to $\mathcal{M}_+$) points in the direction of increasing (resp. decreasing) $r_\pm$. Finally, $R = R(\tau)$ is the radius of the shell (or *bubble*) expressed as a function of the proper time τ of an observer that lives on the bubble itself. Most of the popularity of shell models (particularly in the spherically symmetric version) is likely due to the direct geometrical meaning of the junction conditions, as well as to the fact that in spherical symmetry, it is possible to reduce (1) to

$$\dot{R}^2 + V(R) = 0, \quad V(R) = -\{(R^2 f_-(R) + R^2 f_+(R) - M^2(R))^2 - 4R^4 f_-(R) f_+(R)\} / (4M^2(R) R^2). \quad (2)$$

The solutions of (1) are equivalent to the solutions of (2) when the classical looking equation is complemented by the results $\epsilon_\pm = \text{sign}\{M(R)(R^2 f_-(R) - R^2 f_+(R) \mp M^2(R))\}$, which are required to obtain the global spacetime structure of $\mathcal{S}$ starting from the knowledge of the trajectory $R(\tau)$. Thanks to the fact that the classical dynamics can be exactly solved (at least numerically), it is then tempting to proceed and study its quantum regime [2, 3, 4, 5]. In particular, since it is often the case that $V(R)$ in (2) acts as a potential barrier between bounded and unbounded solutions, it can be interesting to study, both, the semiclassical states corresponding to the bounded solutions [6] as well as the tunnelling under the potential barrier. Both these approaches have been considered, but here we will concentrate only on the second one, i.e. the tunnelling process: while waiting for quantum gravity, the natural framework for a its

¹E-mail: ansoldi@trieste.infn.it — Web-page: <http://www-dft.ts.infn.it/~ansoldi>



complete analysis, many studies have already been performed at the semiclassical level. In particular, for these tunnelling processes we would like to determine: i) the geometry (instanton) interpolating between, for instance, the bounded initial configuration and the unbounded final one (if this instanton exists); ii) a general procedure to calculate the probability for the process. In this contribution we will summarize some problems that appear when trying to implement the above program and which resist, unsolved, since more than fifteen years [7], hoping to shed some light on possible successful approaches. In more detail, the problem of determining the Euclidean solution mediating the tunnelling can be formulated in the framework that we briefly depicted above. The Euclidean junction can be proved to be described by an equation very similar to (1); formally it can be obtained by simply Wick rotating the Euclidean time τ : $\tau \rightarrow \tau^{(E)} = -i\tau$ and correspondingly, $t_{\pm}^{(E)} = -it_{\pm}$. This gives the Wick rotated equation

$$R \left(\epsilon_+ \sqrt{f_+(R) - (R')^2} - \epsilon_- \sqrt{f_-(R) - (R')^2} \right) = M(R) \quad \Rightarrow \quad (R')^2 - V(R) = 0, \quad (3)$$

where a prime denotes a derivative with respect to $\tau^{(E)}$. Under Wick rotation, the results for $\epsilon_{\pm}$ are clearly unchanged. It is also worth remembering the relations $(t_{\pm}^{(E)})' = \epsilon_{\pm} \sqrt{f_{\pm}(R) - (R')^2} / f_{\pm}(R)$, as well as the fact that it is possible to provide an effective Lagrangian formulation [8, 9] to derive (1) and (3). Incidentally, we observe that (1) and (3) are, in fact, first order equations: they are a first integral of the second order Euler-Lagrange equations obtained from the effective Lagrangian L_{eff} . From L_{eff} the effective momentum P_{eff} can be derived with standard techniques and its Euclidean counterpart is

$$P_{\text{eff}}^{(E)} = -R \left\{ \arctan \left(\frac{R'}{\epsilon_+ \sqrt{f_+(R) - (R')^2}} \right) - \arctan \left(\frac{R'}{\epsilon_- \sqrt{f_-(R) - (R')^2}} \right) \right\}. \quad (4)$$

We have, now, at least two choices to determine the transition amplitude: use the path-integral approach with the Lagrangian L_{eff} or proceed *via* canonical quantization using the Euclidean momentum $P_{\text{eff}}^{(E)}$ and the standard result for the probability amplitude $\mathcal{A}$

$$\mathcal{A} \sim \exp(-S), \quad S = \int_{R_1}^{R_2} P(R) dR, \quad R_1, R_2 \text{ extrema of the tunneling trajectory.} \quad (5)$$

For concreteness, let us now specialize to the case considered in [10] where this analysis has been performed by choosing $f_+(r_+) = 1 - \chi^2 r_+^2$, i.e. de Sitter spacetime, $f_-(r_-) = 1 - 2M/r_-$, i.e. Schwarzschild spacetime, and the shell has a constant tension ρ , i.e. $M(R) = 4\pi\rho R^2$. This is a special, important case often considered in the studies of vacuum bubbles/decay [11, 4, 5]. This specific model allows a description of inflation in early universe cosmology avoiding the initial singularity problem. Indeed, it turns out that for a wide range of values of the parameters χ , M and ρ , there are bounded inflating solutions that can be created without an initial singularity; although they do not inflate enough to be a good model of the present universe, it is suggestive to consider the possibility that they will *tunnel* into an unbounded solution which will eventually evolve and resemble the present universe. This idea becomes even more stimulating in view of some issues which appear at a careful analysis of the process [10] and which have not yet found a satisfactory solution/explanation. They are i) the fact that the Euclidean manifold which should mediate the transition between the two Lorentzian junctions (the one before the tunnelling and the one after) can not be easily defined and ii) a discordance between the results provided by a path-integral approach and those obtained with a canonical one. If in the seminal paper of Farhi *et al.* [10] an interesting proposal for the construction of the instanton has been put forward and its generalization to generic junctions might provide the sought answer, the second issue is mostly disturbing in view of the fact that canonical methods are known to reproduce well known results in vacuum decay (in particular the canonical approach has been shown to reproduce [9] the results of Coleman *et al.* [3] and Parke [12]). A slightly more detailed analysis of this point with additional technical details can be found in figure 1. Then, we would like to conclude this contribution with just a few, more speculative, considerations. In particular, what is the physical counterpart of these open issues in the Euclidean sector? Is the tunnelling process really allowed or not? If yes, is there an instanton mediating the process? What is the meaning of the discrepancy between path-integral and canonical formulations? Could it be of interest for quantum gravity? And more, how can we interpret all the above questions in

view of the initial singularity problem? For instance, we could take the point of view that transitions which do not satisfy all the standard properties of the Euclidean momentum are forbidden; but then we would forbid many transitions which would help us to evade classical singularity theorems! If, instead, we can make sense of the unusual properties of the (Euclidean) spacetime structure, we could easily develop a lot of other (solvable) examples in which, already at the semiclassical level, quantum effects can be effectively used to remove singularities. . . . The search for an answer to these and other similar (interesting) questions is, currently, work in progress, and its results will be reported elsewhere.

Acknowledgements

I would like to thank Prof. Gianrossano Giannini for his warm encouragement during the (continuing) development of this project, and Prof. A. Guth, Prof. H. Ishihara and Prof. T. Tanaka for stimulating discussions related to the subject of this contribution. I would also like to gratefully acknowledge partial financial support from the Yukawa Institute of Theoretical Physics, which made possible my participation in the JGRG16 conference.

References

- [1] W. Israel, *Nuovo Cimento B* **XLIV** B:1, 1966; err. ibid. **III**:463(E), 1967; C. Barrabes and W. Israel, *Phys. Rev.* **D43**:1129, 1991.
- [2] S. Coleman, *Phys. Rev.* **D15**:2929, 1977; C. G. Callan Jr. and S. Coleman, *Phys. Rev.* **D16**:1762, 1977; V. A. Berezin, *Phys. Lett. B* **241**:194, 1990; W. Fischler, D. Morgan and J. Polchinski, *Phys. Rev.* **D42**:4042, 1990; M. Visser, *Phys. Rev.* **D43**:402, 1991; M. Visser, *Lorentzian wormholes: from Einstein to Hawking*, American institute of Physics, Woodbury, 1995; K. Nakamura, Y. Oshiro and A. Tomimatsu, *Phys. Rev.* **D53**:4356, 1996; A. D. Dolgov and I. B. Khriplovich, *Phys. Lett. B* **400**:12, 1997; G. L. Alberghi, R. Casadio, G. P. Vacca and G. Venturi, *Class. Quantum Grav.* **16**:131, 1999; N. Sakai, K. Nakao, H. Ishihara and M. Kobayashi, *Phys. Rev.* **D74**:024026, 2006.
- [3] S. Coleman and F. de Luccia, *Phys. Rev.* **D21**:3305, 1980.
- [4] V. A. Berezin, N. G. Kozimirov, V. A. Kuzmin and I. I. Tkachev, *Phys. Lett. B* **212**:415, 1988.
- [5] W. Fischler, D. Morgan and J. Polchinski, *Phys. Rev.* **D41**:2638, 1990.
- [6] S. Ansoldi, *Class. Quantum Grav.* **19**:6231, 2002; S. Ansoldi, *AIP Conf. Proc.* **751**:159, 2005; S. Ansoldi, “Braneworlds and quantum states of relativistic shells”, *To appear in the proceedings of the 11th Marcell Grossman Meeting on General Relativity*, Berlin, July 23-29, 2006; S. Ansoldi, E. I. Guendelman, and H. Ishihara, In preparation.
- [7] A. Aguirre and M. C. Johnson, *Phys. Rev.* **D72**:103525, 2005; A. Aguirre and M. C. Johnson, *Phys. Rev.* **D73**:123529, 2006; S. Ansoldi and L. Sindoni, “Gravitational tunnelling of relativistic shells”, in *Frontiers in Fundamental Physics*, Springer, 2006.
- [8] J. L. Friedman, J. Louko and S. N. Winters-Hilt, *Phys. Rev.* **D56**:7674, 1997; P. Hajicek and J. Bicak, *Phys. Rev.* **D56**:4706, 1997; P. Hajicek, *Phys. Rev.* **D58**:084005, 1998; P. Hajicek, *Phys. Rev.* **D57**:936, 1998; P. Hajicek and J. Kijowski, *Phys. Rev.* **D57**:914, 1998; err. ibid. **D61**:129901(E), 2000; J. Kijowski, *Acta Phys. Pol. B* **29**:1001, 1998; S. Mukohyama, *Phys. Rev.* **D65**:024028, 2002; V. D. Gladush, *J. Math. Phys.* **42**:2590, 2000; V. D. Gladush, *Gen. Rel. Grav.* **36**:1821, 2004; C. Barrabes and W. Israel, *Phys. Rev.* **D71**:064008, 2005.
- [9] S. Ansoldi, A. Aurilia, R. Balbinot and E. Spallucci, *Physics Essays* **9**:556, 1996; S. Ansoldi, A. Aurilia, R. Balbinot and E. Spallucci, *Class. Quantum Gravity* **14**:2727, 1997.
- [10] E. Farhi, A. H. Guth and J. Guven, *Nucl. Phys. B* **339**:417, 1990; A. H. Guth, *Phys. Scr.* **T36**:237, 1991.
- [11] K. Sato, M. Sasaki, H. Kodama and K. Maeda, *Progr. Theor. Phys.* **65**:1443, 1981; H. Kodama, M. Sasaki, K. Sato and K. Maeda, *Progr. Theor. Phys.* **66**:2052, 1981; K. Sato, H. Kodama, M. Sasaki and K. Maeda, *Phys. Lett. B* **108**:103, 1982; H. Kodama, M. Sasaki and K. Sato, *Progr. Theor. Phys.* **68**:1979, 1982; K. Maeda, K. Sato, M. Sasaki and H. Kodama, *Phys. Lett. B* **108**:98, 1982; S. K. Blau, E. I. Guendelman and A. H. Guth, *Phys. Rev.* **D35**:1747, 1987; V. A. Berezin, V. A. Kuzmin and I. I. Tkachev, *Phys. Rev.* **D36**:2919, 1987; A. Aurilia, M. Palmer and E. Spallucci, *Phys. Rev.* **D40**:2511, 1989; M. Sasaki, T. Tanaka, K. Yamamoto and J. Yokoyama, *Phys. Lett. B* **317**:510, 1993; T. Tanaka, *Nucl. Phys. B* **556**:373, 1999; A. Khvedelidze, G. V. Lavrelashvili and T. Tanaka, *Phys. Rev.* **D62**:083501, 2000.
- [12] S. Parke *Phys. Lett. B* **121**:313, 1983.